

(19) World Intellectual Property Organization
International Bureau



BEST AVAILABLE COPY

(43) International Publication Date
4 January 2001 (04.01.2001)

PCT

(10) International Publication Number
WO 01/01133 A3

(51) International Patent Classification⁷: G01N 33/564

(21) International Application Number: PCT/EP00/06025

(22) International Filing Date: 28 June 2000 (28.06.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
99111975.1 28 June 1999 (28.06.1999) EP

(71) Applicants (for all designated States except US): IMMUNDIAGNOSTIK AG [DE/DE]; Wiesenstrasse 4, D-64625 Bensheim (DE). WIESLAB AB [SE/SE]; Ideon, S-223 70 Lund (SE).

(71) Applicant and

(72) Inventor: PAULSSON, Mats [SE/DE]; Institute for Biochemistry II, Medical Faculty, University of Cologne, Joseph-Stelzmann-Str. 52, D-50931 Cologne (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): SARDY, Miklos

[HU/DE]; Institute for Biochemistry II, Medical Faculty, University of Cologne, Joseph-Stelzmann-Str. 52, D-50931 Cologne (DE). ODENTHAL, Uwe [DE/DE]; Institute for Biochemistry II, Medical Faculty, University of Cologne, Joseph-Stelzmann-Str. 52, D-50931 Cologne (DE). KARPATI, Sarolta [HU/HU]; Department of Dermato-Venerology, Semmelweis University of Medicine, Maria u. 41, H-1085 Budapest (HU). SMYTH, Neil [GB/DE]; Institute for Biochemistry II, Medical Faculty, Joseph-Stelzmann-Str. 52, D-50931 Cologne (DE). AESCHLIMANN, Daniel [CH/GB]; Connective Tissue Biology Laboratories, School of Biosciences, Cardiff University, Museums Ave, Cardiff CF10 3US (GB).


(74) Agents: BENEDUM, Ulrich, Max et al.; Haseltine Lake Partners, Rosenheimer Str. 30, D-81669 Munich (DE).

(81) Designated States (national): AU, CA, DE, DE (utility model), JP, US.

(84) Designated States (regional): European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).

[Continued on next page]

(54) Title: DIAGNOSIS OF GLUTEN SENSITIVE ENTEROPATHY AND OTHER AUTOIMMUNOPATHIES



(57) Abstract: Method for diagnosis of autoimmune diseases of the GSE-type or associated with gluten sensitive enteropathy comprising taking a sample and testing the sample for antibodies against human tissue transglutaminase, tissue-specific transglutaminases, or other transglutaminases. It was found that autoimmune diseases other than coeliac disease can be diagnosed and distinguished in this way, notably, dermatitis herpetiformis Duhring, Crohn's disease, Addison's disease, AI haemolytic anaemia, AI thrombocytopenic purpura, AI thyroid diseases, atrophic gastritis - pernicious anaemia, IgA nephropathy or IgA glomerulonephritis, myasthenia gravis, partial lipodystrophy, polymyositis, primary biliary cirrhosis, primary sclerosing cholangitis, recurrent pericarditis, relapsing polychondritis, rheumatoid arthritis, rheumatism, sarcoidosis, Sjögren's syndrome, SLE, splenic atrophy, type I (insulin dependent) diabetes mellitus, diabetes mellitus of other types, ulcerative colitis, vasculitis (both systemic and cutaneous), vitiligo as well as autoimmune diseases associated with infertility, increased risk of abortion, or reduced foetal growth.

WO 01/01133 A3

WO 01/01133 A3



Published:

— *With international search report.*

(88) Date of publication of the international search report:
19 April 2001

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

BEST AVAILABLE COPY

PATENT COOPERATION TREATY

PCT

REC'D 08 AUG 2001

WIPO PCT

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Article 36 and Rule 70)

5

Applicant's or agent's file reference HL-968.001/UMB	FOR FURTHER ACTION See Notification of Transmittal of International Preliminary Examination Report (Form PCT/IPEA/416)	
International application No. PCT/EP00/06025	International filing date (day/month/year) 28/06/2000	Priority date (day/month/year) 28/06/1999
International Patent Classification (IPC) or national classification and IPC G01N33/564		
Applicant PAULSSON, Mats et al.		

1. This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.



2. This REPORT consists of a total of 5 sheets, including this cover sheet.

- ☒ This report is also accompanied by ANNEXES, i.e. sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT).

These annexes consist of a total of 2 sheets.

3. This report contains indications relating to the following items:

- I ☒ Basis of the report
- II ☐ Priority
- III ☐ Non-establishment of opinion with regard to novelty, inventive step and industrial applicability
- IV ☐ Lack of unity of invention
- V ☒ Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement
- VI ☒ Certain documents cited
- VII ☒ Certain defects in the international application
- VIII ☒ Certain observations on the international application

Date of submission of the demand 26/01/2001	Date of completion of this report 03.08.2001
Name and mailing address of the international preliminary examining authority:  European Patent Office D-80298 Munich Tel. +49 89 2399 - 0 Tx: 523656 epmu d Fax: +49 89 2399 - 4465	Authorized officer Hoesel, H Telephone No. +49 89 2399 8693 

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No. PCT/EP00/06025

I. Basis of the report

1. With regard to the **elements** of the international application (*Replacement sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to this report since they do not contain amendments (Rules 70.16 and 70.17)*):

Description, pages:

1-22 as originally filed

Claims, No.:

1-9 as received on 18/07/2001 with letter of 17/07/2001

Drawings, sheets:

1/12-12/12 as originally filed

2. With regard to the **language**, all the elements marked above were available or furnished to this Authority in the language in which the international application was filed, unless otherwise indicated under this item.

These elements were available or furnished to this Authority in the following language: , which is:

- ☐ the language of a translation furnished for the purposes of the international search (under Rule 23.1(b)).
- ☐ the language of publication of the international application (under Rule 48.3(b)).
- ☐ the language of a translation furnished for the purposes of international preliminary examination (under Rule 55.2 and/or 55.3).

3. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international preliminary examination was carried out on the basis of the sequence listing:

- ☐ contained in the international application in written form.
- ☐ filed together with the international application in computer readable form.
- ☐ furnished subsequently to this Authority in written form.
- ☐ furnished subsequently to this Authority in computer readable form.
- ☐ The statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.
- ☐ The statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished.

4. The amendments have resulted in the cancellation of:

- ☐ the description, pages:
- ☐ the claims, Nos.:

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No. PCT/EP00/06025

☐ the drawings, sheets:

5. ☐ This report has been established as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed (Rule 70.2(c)):

(Any replacement sheet containing such amendments must be referred to under item 1 and annexed to this report.)

6. Additional observations, if necessary:

V. Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement

1. Statement

Novelty (N)	Yes:	Claims	1 - 9
	No:	Claims	
Inventive step (IS)	Yes:	Claims	1 - 9
	No:	Claims	
Industrial applicability (IA)	Yes:	Claims	1 - 9
	No:	Claims	

2. Citations and explanations
see separate sheet

VI. Certain documents cited

1. Certain published documents (Rule 70.10)

and / or

2. Non-written disclosures (Rule 70.9)

see separate sheet

VII. Certain defects in the international application

The following defects in the form or contents of the international application have been noted:
see separate sheet

VIII. Certain observations on the international application

The following observations on the clarity of the claims, description, and drawings or on the question whether the claims are fully supported by the description, are made:
see separate sheet

Reference is made to the following documents:

- D1: SEISSLER ET AL: HORMONE AND METABOLIC RES, vol. 31, June 1999
p. 375-379
D2: SARDY ET AL: CLIN. CHEM. vol. 45, December 1999, p. 2142-2149,

SECTION V:

1. Assays for autoantibodies specific for tissue transglutaminase ("TGc") that are based on the use of human TGc in combination with other transglutaminases as test antigens antigen as presently defined in independent claims 1 and 5 is neither disclosed in nor rendered obvious by the prior art referred to in the international search report.

Consequently, claims 1 - 9 appear to fulfil the requirement of Art. 33(2) and (3) PCT (but please note Section VIII, item 4).

SECTION VI:

2. The examination has been based on the presumption that the priority has been validly claimed. If not, the disclosure of D2 would have to be taken into consideration as relevant prior art.

SECTION VII:

3. Contrary to the requirements of Rule 5.1(a)(ii) PCT, the relevant background art disclosed in the document D1 is not mentioned in the description, nor is this document identified therein.

SECTION VIII:

4. The claims do not meet the requirement of art. 6 PCT for the following reasons.

A major part of the experiments concerns tests using one single antigen only, huTGc, huTGe or gpTGc (see p. 10, line 31 - p. 11, line 20, p. 13, line 35 - p. 15,

line 22, p. 14, line 25 - p. 15, lines 4 - 8, p. 15, lines 18 - p. 19, line 7). These tests do not represent "comparative protein binding tests" in the commonly acknowledged technical meaning (comparison of protein binding to at least two different antigens wherein the binding pattern allows diagnostically significant conclusions). The said tests do not fall within the scope of present claims 1 and 5.

Concerning TGe ELISA, it is indicated that this test is less sensitive than the TGc ELISA and that the presence of TGe inhibits binding of GSE sera to TGc. Particularly the latter indicates that TGe and TGc share common epitopes and moreover suggests that in an TGe ELISA actually anti-TGc autoantibodies are detected.

The description fails to disclose a system that requires testing for a comparative binding pattern of sera to these two TG antigens. Additionally, the application does not provide any substantial evidence for a diagnostically relevant association of different TGe/TGc binding patterns.

The description furthermore fails to provide any evidence concerning the presence and significance of auto-antibodies specific for or detectable by transglutaminases known as TGk, TGx, Factor XIIIa or band 4.2.

Taking into account the applicant's comments, it is noted that the gist of the present application with respect to the prior art of D1 is confined to the findings that (i) anti-TGc-autoantibodies can be used as a marker for dermatitis herpetiformis, and (ii) that these antibodies may be detected (in a less sensitive test) by TGe as test antigen.

In summary, the present claims are basically inconsistent with the actual disclosure and not substantially supported.

CLAIMS:

1. Method for differential diagnosis of autoimmune diseases of the GSE-type or associated with gluten sensitive enteropathy comprising taking a sample and testing the sample for antibodies against human tissue transglutaminase and at least one other transglutaminase molecule selected from FXIIIA, TGk, TGx, TGe and Band 4.2.
2. Method as claimed in claim 1, wherein the autoimmune disease is dermatitis herpetiformis, morbus Duhring.
3. Method as claimed in claim 1, wherein the autoimmune disease is selected from Addison's disease, AI haemolytic anaemia, AI thrombocytopenic purpura, AI thyroid diseases, atrophic gastritis - pernicious anaemia, Crohn's disease, colitis ulcerosa, Goodpasture syndrome, IgA nephropathy or IgA glomerulonephritis, myasthenia gravis, partial lipodystrophy, polymyositis, primary biliary cirrhosis, primary sclerosing cholangitis, progressive systemic sclerosis, recurrent pericarditis, relapsing polychondritis, rheumatoid arthritis, rheumatism, sarcoidosis, Sjögren's syndrome, SLE, splenic atrophy, type I (insulin-dependent) diabetes mellitus, diabetes mellitus of other types, Wegener granulomatosis, ulcerative colitis, vasculitis (both systemic and cutaneous), vitiligo.
4. Method as claimed in claim 1, wherein the autoimmune disease is associated with infertility, increased risk of abortion, reduced fetal growth.
5. Comparative protein binding assay for differential diagnosis of autoimmune diseases comprising the

detection of antibodies against transglutaminase, characterised in that the protein binding assay comprises at least two different transglutaminase antigens selected from tissue transglutaminase (TGc), TGk, TGe, TGx, FXIIIA and Band 4.2

6. Comparative protein binding assay as claimed in claim 5, comprising transglutaminase from different species.

7. Comparative protein binding assay as claimed in any preceding claim 5 to 6, wherein the transglutaminase antigen is a recombinant fusion protein or fragment thereof.

8. Comparative protein binding assay as claimed in any preceding claim 5 to 7, wherein the binding assay is an immunoassay selected from RIA, EIA/ELISA, LiA and FIA.

9. Comparative protein binding assay as claimed in any preceding claim 5 to 8, wherein the binding assay is a sandwich-immunoassay selected from IRMA, IEMA/EUA, ILMA (immunoluminescence assay) and IFMA (immuno-fluorescence assay).

CLAIMS:

- 5 1. Method for diagnosis of autoimmune diseases of the GSE-type or associated with gluten sensitive enteropathy comprising taking a sample and testing the sample for antibodies against human tissue transglutaminase and at least one other transglutaminase molecule.
- 10 2. Method as claimed in claim 1, wherein the autoimmune disease is dermatitis herpetiformis, morbus Duhring.
- 15 3. Method as claimed in claim 1, wherein the autoimmune disease is selected from Addison's disease, AI haemolytic anaemia, AI thrombocytopenic purpura, AI thyroid diseases, atrophic gastritis - pernicious anaemia, Crohn's disease, colitis ulcerosa, Goodpasture syndrome, IgA nephropathy or IgA glomerulonephritis, myasthenia gravis, partial lipodystrophy, polymyositis, primary biliary cirrhosis, primary sclerosing cholangitis, progressive systemic sclerosis, recurrent pericarditis, relapsing polychondritis, rheumatoid arthritis, rheumatism, sarcoidosis, Sjögren's syndrome, SLE, splenic atrophy, type I (insulin-dependent) diabetes mellitus, diabetes mellitus of other types, Wegener granulomatosis, ulcerative colitis, vasculitis (both systemic and cutaneous), vitiligo.
- 20 4. Method as claimed in claim 1, wherein the autoimmune disease is associated with infertility, increased risk of abortion, reduced foetal growth.
- 25 5. Comparative protein binding assay for a differential diagnosis of autoimmune diseases comprising the detection of antibodies against transglutaminase, characterised in that the protein binding assay comprises recombinant human tissue transglutaminase as antigen.
- 30 6. Comparative protein binding assay as claimed in claim 5, further comprising tissue-specific transglutaminase as antigen.
7. Comparative protein binding assay as claimed in claim 6, wherein the transglutaminase is selected from FXIIIA, TGk, TGe, TGx, and band 4.2.
- 35 8. Comparative protein binding assay as claimed in any preceding claim 5 or 6, further comprising any other transglutaminase as antigen.

9. Comparativ protein binding assay as claimed in claim 8, comprising transglutaminase from different species.
10. Comparative protein binding assay as claimed in claim 8, wherein the transglutaminase is a
5 recombinant fusion protein or fragment thereof.
11. Comparative protein binding assay as claimed in any preceding claim 5 to 10, wherein the binding assay is an immunoassay selected from RIA, EIA/ELISA, LiA and FiA.
- 10 12. Comparative protein binding assay as claimed in any preceding claim 5 to 11, wherein the binding assay is a sandwich-immunoassay selected from IRMA, IEMA/EUA, ILMA (immunoluminescence assay) and IFMA (immunofluorescence assay).

PATENT COOPERATION TREATY

PCT

INTERNATIONAL SEARCH REPORT

(PCT Article 18 and Rules 43 and 44)

Applicant's or agent's file reference HL-968.001/UMB	FOR FURTHER ACTION see Notification of Transmittal of International Search Report (Form PCT/ISA/220) as well as, where applicable, item 5 below.	
International application No. PCT/EP 00/06025	International filing date (day/month/year) 28/06/2000	(Earliest) Priority Date (day/month/year) 28/06/1999
Applicant PAULSSON, Mats et al.		

This International Search Report has been prepared by this International Searching Authority and is transmitted to the applicant according to Article 18. A copy is being transmitted to the International Bureau.

This International Search Report consists of a total of 3 sheets.



It is also accompanied by a copy of each prior art document cited in this report.

1. Basis of the report

- a. With regard to the **language**, the international search was carried out on the basis of the international application in the language in which it was filed, unless otherwise indicated under this item.



the international search was carried out on the basis of a translation of the international application furnished to this Authority (Rule 23.1(b)).

- b. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international search was carried out on the basis of the sequence listing:



contained in the international application in written form.



filed together with the international application in computer readable form.



furnished subsequently to this Authority in written form.



furnished subsequently to this Authority in computer readable form.



the statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.



the statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished

2. ☐ **Certain claims were found unsearchable** (See Box I).

3. ☐ **Unity of invention is lacking** (see Box II).

4. With regard to the **title**,



the text is approved as submitted by the applicant.



the text has been established by this Authority to read as follows:

5. With regard to the **abstract**,



the text is approved as submitted by the applicant.



the text has been established, according to Rule 38.2(b), by this Authority as it appears in Box III. The applicant may, within one month from the date of mailing of this international search report, submit comments to this Authority.

6. The figure of the **drawings** to be published with the abstract is Figure No.



as suggested by the applicant.



because the applicant failed to suggest a figure.



because this figure better characterizes the invention.

1



None of the figures.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 00/06025

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 G01N33/564

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 G01N C12N C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	SARDY MIKLOS ET AL: "Recombinant human tissue transglutaminase ELISA for the diagnosis of gluten-sensitive enteropathy." CLINICAL CHEMISTRY, vol. 45, no. 12, December 1999 (1999-12), pages 2142-2149, XP002155800 ISSN: 0009-9147 the whole document --- -/--	1-12

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

G document member of the same patent family

Date of the actual completion of the international search

3 January 2001

Date of mailing of the international search report

19/01/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Pellegrini, P

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 00/06025

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	SEISSLER J ET AL: "Antibodies to human recombinant tissue transglutaminase measured by radioligand assay: Evidence for high diagnostic sensitivity for celiac disease." HORMONE AND METABOLIC RESEARCH, vol. 31, no. 6, June 1999 (1999-06), pages 375-379, XP000974671 ISSN: 0018-5043 the whole document ----	1-12
X	DE 196 30 557 A (DIETERICH WALBURGA ;SCHUPPAN DETLEF PRIV DOZ DR DR (DE)) 29 January 1998 (1998-01-29) the whole document ----	1-12
X	DIETERICH WALBURGA ET AL: "Identification of tissue transglutaminase as the autoantigen of celiac disease." NATURE MEDICINE, vol. 3, no. 7, 1997, pages 797-801, XP002155802 ISSN: 1078-8956 cited in the application the whole document ----	1-12
A	US 5 726 051 A (BIRCKBICHLER PAUL J ET AL) 10 March 1998 (1998-03-10) claims -----	1-12

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 00/06025

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 19630557 A	29-01-1998	AU 718797 B	20-04-2000
		AU 4201197 A	10-02-1998
		BR 9710500 A	18-01-2000
		CN 1225723 A	11-08-1999
		CZ 9900117 A	14-07-1999
		WO 9803872 A	29-01-1998
		EP 0912898 A	06-05-1999
		ES 2131038 T	16-07-1999
		GR 99300020 T	30-06-1999
		NO 990190 A	15-03-1999
		PL 331203 A	05-07-1999
		SI 9720044 A	31-08-1999
		SK 6799 A	16-05-2000
US 5726051 A	10-03-1998	AU 5590394 A	24-05-1994
		WO 9410296 A	11-05-1994

PATENT COOPERATION TREATY

PCT

INTERNATIONAL SEARCH REPORT

(PCT Article 18 and Rules 43 and 44)

Applicant's or agent's file reference HL-968.001/UMB	FOR FURTHER ACTION see Notification of Transmittal of International Search Report (Form PCT/ISA/220) as well as, where applicable, item 5 below.	
International application No. PCT/EP 00/ 06025	International filing date (<i>day/month/year</i>) 28/06/2000	(Earliest) Priority Date (<i>day/month/year</i>) 28/06/1999
Applicant PAULSSON, Mats et al.		

This International Search Report has been prepared by this International Searching Authority and is transmitted to the applicant according to Article 18. A copy is being transmitted to the International Bureau.

This International Search Report consists of a total of 3 sheets.
☒ It is also accompanied by a copy of each prior art document cited in this report.

1. Basis of the report

- a. With regard to the **language**, the international search was carried out on the basis of the international application in the language in which it was filed, unless otherwise indicated under this item.
- ☐ the international search was carried out on the basis of a translation of the international application furnished to this Authority (Rule 23.1(b)).
- b. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international search was carried out on the basis of the sequence listing :
- ☐ contained in the international application in written form.
- ☐ filed together with the international application in computer readable form.
- ☐ furnished subsequently to this Authority in written form.
- ☐ furnished subsequently to this Authority in computer readable form.
- ☐ the statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.
- ☐ the statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished.

2. ☐ **Certain claims were found unsearchable** (See Box I).

3. ☐ **Unity of invention is lacking** (see Box II).

4. With regard to the **title**,

- ☒ the text is approved as submitted by the applicant.
- ☐ the text has been established by this Authority to read as follows:

5. With regard to the **abstract**,

- ☒ the text is approved as submitted by the applicant.
- ☐ the text has been established, according to Rule 38.2(b), by this Authority as it appears in Box III. The applicant may, within one month from the date of mailing of this international search report, submit comments to this Authority.

6. The figure of the **drawings** to be published with the abstract is Figure No.

- ☐ as suggested by the applicant.
- ☒ because the applicant failed to suggest a figure.
- ☐ because this figure better characterizes the invention.

1
☐ None of the figures.

INTERNATIONAL SEARCH REPORT
Information on patent family members

Inter Application No
PCT/EP 00/06025

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 19630557 A	29-01-1998	AU 718797 B	20-04-2000
		AU 4201197 A	10-02-1998
		BR 9710500 A	18-01-2000
		CN 1225723 A	11-08-1999
		CZ 9900117 A	14-07-1999
		WO 9803872 A	29-01-1998
		EP 0912898 A	06-05-1999
		ES 2131038 T	16-07-1999
		GR 99300020 T	30-06-1999
		NO 990190 A	15-03-1999
		PL 331203 A	05-07-1999
		SI 9720044 A	31-08-1999
		SK 6799 A	16-05-2000
US 5726051 A	10-03-1998	AU 5590394 A	24-05-1994
		WO 9410296 A	11-05-1994

BEST AVAILABLE COPY

INTERNATIONAL SEARCH REPORT

Int'l Application No

PCT/EP 00/06025

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 601N33/564

BEST AVAILABLE COPY

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 601N C12N C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	SARDY MIKLOS ET AL: "Recombinant human tissue transglutaminase ELISA for the diagnosis of gluten-sensitive enteropathy." CLINICAL CHEMISTRY, vol. 45, no. 12, December 1999 (1999-12), pages 2142-2149, XP002155800 ISSN: 0009-9147 the whole document --- -/--	1-12

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

13 January 2001

Date of mailing of the international search report

19/01/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Pellegrini, P

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	SEISSLER J ET AL: "Antibodies to human recombinant tissue transglutaminase measured by radioligand assay: Evidence for high diagnostic sensitivity for celiac disease." HORMONE AND METABOLIC RESEARCH, vol. 31, no. 6, June 1999 (1999-06), pages 375-379, XP000974671 ISSN: 0018-5043 the whole document	1-12
X	DE 196 30 557 A (DIETERICH WALBURGA ;SCHUPPAN DETLEF PRIV DOZ DR DR (DE)) 29 January 1998 (1998-01-29) the whole document	1-12
X	DIETERICH WALBURGA ET AL: "Identification of tissue transglutaminase as the autoantigen of celiac disease." NATURE MEDICINE, vol. 3, no. 7, 1997, pages 797-801, XP002155802 ISSN: 1078-8956 cited in the application the whole document	1-12
A	US 5 726 051 A (BIRCKBICHLER PAUL J ET AL) 10 March 1998 (1998-03-10) claims	1-12

BEST AVAILABLE COPY

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
4 January 2001 (04.01.2001)

PCT

(10) International Publication Number
WO 01/01133 A2

(51) International Patent Classification⁷: G01N 33/53

(21) International Application Number: PCT/EP00/06025

(22) International Filing Date: 28 June 2000 (28.06.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
99111975.1 28 June 1999 (28.06.1999) EP

(71) Applicants (*for all designated States except US*): IM-
MUNDIAGNOSTIK AG [DE/DE]; Wiesenstrasse 4,
D-64625 Bensheim (DE). WIESLAB AB [SE/SE]; Ideon,
S-223 70 Lund (SE).

(71) Applicant and

(72) Inventor: PAULSSON, Mats [SE/DE]; Institute for
Biochemistry II, Medical Faculty, University of Cologne,
Joseph-Stelzmann-Str. 52, D-50931 Cologne (DE).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): SARDY, Miklos
[HU/DE]; Institute for Biochemistry II, Medical Fac-
ulty, University of Cologne, Joseph-Stelzmann-Str. 52,
D-50931 Cologne (DE). ODENTHAL, Uwe [DE/DE];

Institute for Biochemistry II, Medical Faculty, University
of Cologne, Joseph-Stelzmann-Str. 52, D-50931 Cologne
(DE). KARPATI, Sarolta [HU/HU]; Department of Der-
mato-Venerology, Semmelweis University of Medicine,
Maria u. 41, H-1085 Budapest (HU). SMYTH, Neil
[GB/DE]; Institute for Biochemistry II, Medical Fac-
ulty, Joseph-Stelzmann-Str. 52, D-50931 Cologne (DE).
AESCHLIMANN, Daniel [CH/GB]; Connective Tissue
Biology Laboratories, School of Biosciences, Cardiff
University, Museums Ave, Cardiff CF10 3US (GB).

(74) Agents: BENEDUM, Ulrich, Max et al.; Haseltine Lake
Partners, Rosenheimer Str. 30, D-81669 Munich (DE).

(81) Designated States (*national*): AU, CA, DE, DE (utility
model), JP, US.

(84) Designated States (*regional*): European patent (AT, BE,
CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC,
NL, PT, SE).

Published:

— *Without international search report and to be republished
upon receipt of that report.*

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: DIAGNOSIS OF GLUTEN SENSITIVE ENTEROPATHY AND OTHER AUTOIMMUNOPATHIES

(57) Abstract: Method for diagnosis of autoimmune diseases of the GSE-type or associated with gluten sensitive enteropathy comprising taking a sample and testing the sample for antibodies against human tissue transglutaminase, tissue-specific transglutaminases, or other transglutaminases. It was found that autoimmune diseases other than coeliac disease can be diagnosed and distinguished in this way, notably, dermatitis herpetiformis, Dühring, Crohn's disease, Addison's disease, AI haemolytic anaemia, AI thrombocytopenic purpura, AI thyroid diseases, atrophic gastritis - pernicious anaemia, IgA nephropathy or IgA glomerulonephritis, myasthenia gravis, partial lipodystrophy, polymyositis, primary biliary cirrhosis, primary sclerosing cholangitis, recurrent pericarditis, relapsing polychondritis, rheumatoid arthritis, rheumatism, sarcoidosis, Sjögren's syndrome, SLE, splenic atrophy, type I (insulin-dependent) diabetes mellitus, diabetes mellitus of other types, ulcerative colitis, vasculitis (both systemic and cutaneous), vitiligo as well as autoimmune diseases associated with infertility, increased risk of abortion, or reduced foetal growth.

WO 01/01133 A2



Published:

— *With international search report.*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(88) Date of publication of the international search report:

19 April 2001

DIAGNOSIS OF GLUTEN SENSITIVE ENTEROPATHY AND OTHER AUTOIMMUNOPATHIES

Technical Field

5 The invention relates to the diagnosis of gluten sensitive enteropathy and other autoimmune diseases comprising the detection of antibodies against transglutaminases.

Background Art

10 Gluten sensitive enteropathy (GSE) is a common chronic small bowel disorder of autoimmune origin occurring both in children and adults. It is evoked and maintained by wheat gluten which is also present in other cereals. The clinical appearance of GSE is typically coeliac disease (CD). In some individuals, however, it is associated with dermatitis herpetiformis (DH), a bullous, autoimmune skin disease characterised by granular IgA staining of the papillary dermis. Both of these forms of GSE have the same genetic background and have been associated with HLA class II antigens DQ2, and DR3, and the HLA-A1, -B8, -DR3 haplotype. In case of ingestion of gluten and the
15 presence of genetic predisposition, a T-cell mediated autoimmune response develops in the small bowel first resulting in lymphocytic infiltration, later reduction followed by total atrophy of the villi leading to a disturbed resorption (Marsh M N et al., Bailliere Clin Gastr 1995;9:273-294). On a completely gluten-free diet, however, the pathological alterations entirely disappear and a normal morphology and function is restored.

20

The diagnosis of GSE is based on the characteristic histological changes (villous atrophy, intraepithelial lymphocytosis, crypt hyperplasia) seen in jejunal biopsies, followed by the regeneration of the mucosa after a gluten-free diet and relapse during subsequent gluten challenge. However, serological tests may be helpful in the diagnosis of GSE as they offer a less invasive and cheaper
25 alternative. These detect IgA antibodies directed against endomysial antigen, reticulin, or gliadin. The IgA-class endomysial antibody (EMA) test is considered to be the serological method of choice, because of its higher sensitivity and specificity when compared to the IgA-class anti-reticulin antibody and the IgA-class anti-gliadin antibody tests. EMA is found in 60-70% of untreated patients with DH and in almost all untreated patients with CD. The EMA test is performed, however, on expensive
30 oesophagus sections from endangered primates, is laborious and time consuming, and subjective in borderline cases.

Tissue transglutaminase (TGc, EC 2.3.2.13) was further identified as the predominant or sole endomysial autoantigen of CD (Dieterich W et al., Nature Medicine 1997, 3(7):797-801). An ELISA
35 test for CD has been produced based upon the commercially available guinea pig TGc. Although the amino acid sequence identity between guinea pig and human TGc's is 82.8%, this test gave high sensitivity and specificity above 90% (Ikura K et al., Biochemistry 1988;27:2898-905; Gentile V et al., J

Biol Chem 1991;266:478-83; Dieterich W et al., Gastroenterology 1998;115:1317-21; Sulkanen S et al, Gastroenterology 1998;115:1322-8).

Object of Invention

5 It was an object of the invention to provide an improved assay for gluten sensitive enteropathies and in particular an antibody binding assay which allows a differential diagnosis of autoimmune diseases of the GSE-type, autoimmune diseases associated with GSE and seemingly non-active, latent gluten sensitive enteropathies.

10 Brief Description of Invention

This object has been achieved by providing a comparative protein binding assay based on recombinant human tissue transglutaminase and other transglutaminases for a differential diagnosis of gluten sensitive enteropathy (GSE), autoimmune diseases of the GSE-type, autoimmune diseases associated with GSE and a group of autoimmune diseases associated with IgA and IgG against trans-
15 glutaminases. A preferred embodiment of the invention relates to a multiple protein binding assay comprising human tissue transglutaminase and other transglutaminases as binding partners. A more preferred embodiment of the invention relates to a protein binding assay comprising transglutaminases isolated or cloned from different tissues and species.

20 Another aspect of the invention relates to a method for diagnosis of autoimmune diseases of the GSE-type comprising a multiple protein binding assay on the basis of human TGc and other transglutaminases such as guinea pig TGc or other members of the transglutaminase protein family. Multiple or comparative protein binding assay means herewith that the diagnosis is done on basis of at least two, preferably three or more differing transglutaminase molecules as antigens.

25 The invention and preferred modifications thereof are disclosed in claims 1 and 5. Preferred embodiments thereof are described in the dependent claims.

The present invention provide a method for diagnosis of autoimmune diseases of the GSE-
30 type or associated with gluten sensitive enteropathy comprising taking a sample and testing the sample for antibodies against tissue transglutaminase and at least one other transglutaminase. In a preferred embodiment, the autoimmune disease is dermatitis herpetiformis, morbus Duhring, or an autoimmune disease selected from Addison's disease, AI (AI - autoimmune) haemolytic anaemia, AI thrombocytopenic purpura, AI thyroid diseases, IDDM, alopecia, atrophic gastritis - pernicious
35 anaemia, Crohn's disease, hypoadrenalism, hypogonadism, hyposplenism, cryoglobulinism, colitis ulcerosa, Goodpasture syndrome, gluten-induced ataxia, IgA nephropathy or IgA glomerulonephritis, myasthenia gravis, partial lipodystrophy, polymyositis, primary biliary cirrhosis, primary sclerosing

cholangitis, progressive systemic sclerosis, oral aphthosis, recurrent pericarditis, relapsing polychondritis, rheumatoid arthritis, rheumatism, sarcoidosis, sensory neuropathy, seizures, Sjögren's syndrome, SLE, splenic atrophy, type I (insulin-dependent) diabetes mellitus, diabetes mellitus of other types, transaminitis, Wegener granulomatosis, ulcerative colitis, vasculitis (both systemic and cutaneous), vitiligo. A further group of autoimmune diseases that can be diagnosed or distinguished in this way is associated with infertility, increased risk of abortion and/or reduced foetal growth.

The invention also provides a comparative protein binding assay for a differential diagnosis of autoimmune diseases comprising the detection of antibodies against transglutaminase, which protein binding assay comprises recombinant human tissue transglutaminase as antigen. In a preferred embodiment, the comparative protein binding assay comprises further a tissue-specific transglutaminase as antigen, preferably, a transglutaminase selected from FXIIIA, TGk, TGe, TGx, and so-called Band 4.2. The comparative protein binding assay may also contain any other transglutaminase as antigen, in particular, transglutaminase from different species. The skilled person will appreciate that a recombinant fusion protein or fragment thereof can also be used. The comparative protein binding assay is preferably an immunoassay selected from RIA, EIA/ELISA, LiA and FiA and most preferably a sandwich-immunoassay selected from IRMA, IEMA/EUA, ILMA (immunoluminescence assay) and IFMA (immunofluorescence assay).

Astonishingly, it has been found that despite the high amino acid sequence identity between human and guinea pig TGc, some of the tested patient sera not recognised by the guinea pig TGc ELISA had antibodies directed against epitopes of human TGc not conserved in the guinea pig enzyme. Our studies support that TGc is the autoantigen of EMA positive patients also with DH. Moreover, we discovered that there is a class of autoimmune diseases wherein transglutaminases plays a decisive role in the aetiology of the disease, which autoimmune diseases can be distinguished via the unique epitope patterns of the various transglutaminases.

The discovery of TGc as the main autoantigen in GSE did not answer the question why only a proportion of patients with CD also show symptoms of DH and if, whether there is a difference in the antigenic repertoire in these diseases. It was unknown whether the granular IgA precipitates in the skin of DH patients are directed against an antigen present in the skin or if they represent circulating immune complexes deposited in the papillary dermis. The IgA precipitates have so far not been extracted from skin and characterised. However, immunostaining for TGc does not give the same staining pattern as can be seen by direct immunofluorescence studies for IgA precipitates in the skin of DH patients. This suggests that the antigen against which the deposited IgA antibodies are directed might be different from the TGc. We now assume that these antibodies derive from cross-reactivity with another transglutaminase being present in the papillary dermis. Such a limited cross-reactivity

could be a plausible explanation for the moderate penetrance of skin eruptions in patients with CD. It is also unclear why the pathological changes and clinical symptoms have a very limited localisation with only the small intestine and, in case of DH, the skin being affected, as TGc is present in almost every tissue of the human body. To address these questions, we expressed three other
5 transglutaminases which can be found in the skin, the human TGk, TGe and TGx, in human embryonic kidney cells, to measure circulating IgA and IgG titres in patients with CD, DH and other autoimmune diseases.

Brief Description of Drawings

10 The invention will now be described in detail with reference to the accompanying drawings and representations. No limitation of the invention whatsoever shall be construed from the description of the representative examples.

15 **Fig. 1** SDS-PAGE (A) and immunoblot (B) analysis of TGc. The immunoblot was performed using monoclonal antibodies against TGc. Positions of molecular mass standards (kDa) are indicated on the left. Lane I, guinea pig TGc; lane II, lysate of cells producing recombinant human TGc before purification; lane III, flow through; lane IV, eluted TGc from the column.

20 **Fig. 2** Receiver operating characteristic curve (ROC) for the human TGc ELISA showing the point of greatest efficiency of the test, upon which the cut-off level was chosen.

25 **Fig. 3** Serum anti-TGc antibody concentrations in the human TGc ELISA system in the controls (I) and in the patients having CD or DH (II). Treated CD or DH patients are indicated by empty squares. The chosen arbitrary cut-off level for positivity (broken line) is drawn at the AU of 18.

Fig. 4 ROC curve for the guinea pig TGc ELISA showing the point of greatest efficiency of the test, upon which the cut-off level was chosen.

30 **Fig. 5** Serum anti-TGc antibody concentrations in the guinea pig TGc ELISA system in the controls (I) and in the patients having CD or DH (II). Treated CD or DH patients are indicated by empty squares. The chosen arbitrary cut-off level for positivity (broken line) is drawn at the AU of 14.

Fig. 6 Serum concentrations of IgA antibodies against TGc in the human TGc ELISA given in arbitrary units (AU). Treated CD or DH patients are indicated by empty squares. The chosen arbitrary cut-off for positivity (dashed line) is drawn at 18 AU. CD, coeliac disease; DH, dermatitis herpetiformis; CTRL, control sera; PV, pemphigus vulgaris; BP, bullous pemphigoid; CR, Crohn disease; CU, colitis ulcerosa; GP, Goodpasture syndrome; WEG, Wegener granulomatosis; RA, rheumatoid arthritis; SLE, systemic lupus erythematosus; PSS, progressive systemic sclerosis; PA, psoriatic arthritis; HEPC, hepatitis C.

Fig. 7 Box and whisker diagram showing the serum concentrations of IgA antibodies against human TGc. The lower and upper edges of the boxes represent the 25% and 75% percentiles, respectively. The median is indicated by a horizontal line through the box. The lower and upper whiskers represent the 5% and 95% percentiles, respectively. Only untreated patients are included in the CD and DH groups. For abbreviations see Fig. 6.

Fig. 8 Correlation of antibody titres between the human and the guinea pig TGc ELISA. On the x-axis are titres given in arbitrary units measured by the human TGc ELISA, on the y-axis are those measured by the guinea pig TGc ELISA. The cut-off values are presented by dashed lines.

Fig. 9 SDS-PAGE analysis of TGe after purification. Positions of molecular mass standards (kDa) are indicated on the left.

Fig. 10 Immunoblot analysis of the human TGs using polyclonal rabbit antibodies against the Strep II tag. Positions of molecular mass standards (kDa) are indicated on the left. Lane I, TGc; lane II, TGe; lane III, TGk; lane IV, TGx.

Fig. 11 Binding of IgA in GSE patient sera to human TGc or TGe in ELISA. Sera reacting with the TGc showing values below 0.4 gave similar results with the TGe, but sera with elevated IgA antibody titres against TGc in the most cases showed lower antibody titres against TGe.

Fig. 12 Inhibition of binding of IgA in four GSE patient sera to solid phase human TGc by preincubation with increasing amounts of human TGc (dotted line) or TGe (continuous line). Both the TGc and the TGe showed an inhibitory effect, but the TGc was more effective than the TGe.

Detailed Description of Invention

The inventors expressed human TGc and other transglutaminases by recombinant DNA methods and set up various ELISAs based on the purified proteins for detecting IgA anti-TG antibodies. The results of these assays were compared to those from the ELISA with the guinea pig TGc and the EMA test on monkey oesophagus.

Methods and Materials

SDS-PAGE and Immunoblotting

SDS-PAGE (*Sodium dodecyl sulphate polyacrylamide gel electrophoresis*) was performed according to the method of Laemmli (Laemmli, Nature 1970;227:680-5) using a 12% polyacrylamide separating gel with a 5% polyacrylamide stacking gel. Samples were reduced by addition of 2% (v/v) 2-mercaptoethanol. Proteins were detected either by staining with Coomassie Brilliant Blue R (Serva) or by immunoblotting after electrophoretic transfer to a nitro-cellulose membrane (Protran, Schleicher & Schuell) according to the method of Towbin and co-workers (Towbin H et al, Proc Natl Acad Sci USA, 1979;76:4350-4). After protein transfer, the membranes were stained with Ponceau S (Serva), then blocked with 50 mM Tris, 150 mM NaCl, pH 7.4 (TBS) containing 5% non-fat milk powder for 75 min. at room temperature. The blocked membrane was then incubated with the specific antibody.

Human TGc was detected with mouse monoclonal antibodies against TGc (specific for TGc, but cross-reacting both with human and guinea pig TGc, Neomarkers, Ab-3, CUB7402 + TG100) diluted 1:2000 in TBS containing 5% non-fat milk powder and 0.05% Tween™ 20 (Sigma) for 1.5 hours at room temperature. For detection of bound mouse antibodies, membranes were incubated with horse radish peroxidase labelled rabbit antibodies directed against mouse immunoglobulins (Dako), diluted 1:2000 in TBS/Tween containing 5% non-fat milk powder for 1 hour at room temperature. Bound secondary antibodies were detected by autoradiography using the enhanced chemiluminescence system (ECL Kit, Amersham). Guinea pig TGc (Sigma) was used each time as a positive control.

Recombinantly expressed TGs were detected by using rabbit polyclonal antibodies against the Strep II tag (Institut für Bioanalytik) diluted 1:5000 in TBS containing 1% non-fat milk powder and 0.05% Tween 20 (Sigma) for 1 h at room temperature. For detection of bound rabbit antibodies, membranes were incubated with horseradish-peroxidase-labelled swine antibodies against rabbit immunoglobulins (Dako), diluted 1:3000 in TBS/Tween containing 1% non-fat milk powder for 1 hour at room temperature. Bound secondary antibodies were detected by luminescence after incubation 5 min in 100 mM Tris/HCl, pH 8.3, containing 0.2 mM p-coumaric acid (Sigma), 2.65 mM H₂O₂ (Sigma) and 1.25 mM 3-aminophthalhydrazide (Fluka). Human TGc expressed as a fusion protein with a C-terminal Strep II tag was used as a positive control.

Reverse transcription and polymerase chain reaction.

Reverse transcription was performed using AMV reverse transcriptase (Oncor Appligene) and PCR using Pfx Taq polymerase (Life Technologies) following the protocols provided by the manufacturers.

5

Recombinant expression of human TGc.

The episomal eukaryotic expression vector pCEP-Pu/BM40SP, produced from pCEP4 (Invitrogen) (Kohfeldt E et al, FEBS Lett 1997;414:557-61), was modified to introduce a sequence encoding the Strep II tag (IBA, Germany) and a stop codon into the multiple cloning site. The primers
10 5'-GGCCGCATGGAGCCATCCACAATTCGAAAAGTA and 5'-GGCCTACTTTTCGAATTGTGGATG-GCTCCATGC were annealed together and introduced into the Not I site thus constructing a vector (pCEP-Pu/BM40SP/C-Strep) producing a carboxyterminal Strep II fusion protein suitable for streptavidin affinity purification by a StrepTactin™ (IBA, Germany) affinity column as described before by Schmidt TGM et al (J Mol Biol 1996;255:753-66). The full-length human TGc cDNA (GeneBank
15 accession number M55153, cloned in pSP73) was amplified by polymerase chain reaction (PCR) using the 5'-primer 5'-ATTAAGCTTGCCGCCACCATGGCCGAGGAGCTGGTC, and the 3'-primer 5'-TAAGCGGCCGCGGGGCCAATGATGACATTC. The 5'-primer introduced a new Hind III restriction site and a Kozak's translation initiation sequence, the 3'-primer inserted a new Not I restriction site and removed the stop codon. The Hind III/Not I restriction enzyme digested PCR
20 product was purified and inserted at the same restriction sites of the pCEP-Pu/BM40SP/C-Strep, in order to obtain the final expression vector pCEP-Pu/TGc/C-Strep. The correct insertion and sequence of the full construct was verified by cycle sequencing using the ABI Prism Big Dye Terminator Cycle Sequencing Ready Reaction Kit and the products were resolved on an ABI Prism 377 Automated Sequencer (Perkin-Elmer/Applied Biosystems).

25 Human embryonic kidney cells (293-EBNA, Invitrogen) were transfected with pCEP-Pu/TGc/C-Strep and harvested in cell culture in Dulbecco's MEM NUT MIX F-12 (Life Technologies) medium containing 10% foetal bovine serum (Life Technologies), 1% L-glutamine (Life Technologies), 200 IU/ml penicillin (Life Technologies), and 200 µg/ml streptomycin (Life Technologies). Cells were selected with 0.5 µg/ml puromycin. After removing the medium and
30 washing with cold (4°C) 0.25 M sucrose, the cells were lysed mechanically in cold 0.25 M sucrose. The lysate was cleared of particulate material by centrifugation at 27,200 g for 30 min at 4°C, followed by ultracentrifugation of the supernatant at 210,000 g for 60 min at 4°C. After filtering with cheesecloth and adding 1 mM phenylmethanesulfonyl fluoride (PMSF, Fluka) as proteinase inhibitor, 12 ml of the supernatant was passed over a StrepTactin affinity column of 3 cm³ volume equilibrated with sterile
35 and filtered 50 mM Tris/HCl, pH 7.5, containing 1 mM ethylenediamine tetraacetate (EDTA), at 4°C, and at a flow rate of 0.4 ml/cm²/min. After extensive washing with equilibration buffer containing 1 mM PMSF, at a flow rate of 0.9 ml/cm²/min, the protein was eluted with equilibration buffer containing 1

mM PMSF and 2.5 mM desthiobiotin (Sigma), at a flow rate of 0.4 ml/cm²/min. 2 ml fractions were collected. The purification was controlled by Coomassie-stained SDS-PAGE and Western Blotting with monoclonal antibodies against TGc as described above. The protein concentration was estimated by SDS-PAGE, and measured using the bicinchoninic acid (BCA) protein assay reagent (Pierce) following the protocol provided by the supplier, with bovine serum albumin as a standard.

Production of cDNAs from various transglutaminases.

Total RNA from human keratinocytes was reversely transcribed with primers specific for the TGe and TGk proenzymes, and PCR used to obtain cDNAs. Full-length human TGx cDNA (GeneBank accession number AF035960) was cloned in pSP73.

Production of an expression construct for the expression of TGe.

An episomal expression construct was built allowing the production of a C-terminal Strep II fusion protein suitable for streptavidin affinity purification by a StrepTactinTM (Institut für Bioanalytik, Germany) affinity column (Schmidt TGM et al., J Mol Biol 1996;255:753-66.). The TGe proenzyme cDNA was amplified by PCR using the 5'-primer 5'-ATTAAGCTTGCCGCCACCATGGCTGCTCTA-GGAGTC, and the 3'-primer 5'-ATTGCGGCCGCTTCGGCTACATCGATGGACAAC. The 5'-primer introduced a new Hind III restriction site and a Kozak's translation initiation sequence, while the 3'-primer inserted a new Not I restriction site and removed the stop codon. The Hind III/Not I restriction enzyme digested PCR product was purified and inserted at the same restriction sites of the episomal eukaryotic expression vector pCEP-Pu/BM40SP/C-Strep, produced from pCEP4 (Invitrogen), in order to obtain the final expression vector pCEP-Pu/TGe/C-Strep.

Construction of the expression vector pCEP4/N-Strep.

The eukaryotic episomal expression vector pCEP4 (Invitrogen) was modified to introduce a sequence encoding a Kozak's translation initiation sequence and the Strep II tag into the multiple cloning site. The primers 5'-CTAGTTGCCGCCACCATGGCTTGGAGCCATCCACAATTCGAAAAGG and 5'-CTAGCGCCTTTTCGAATTGTGGATGGCTCCAAGCCATGGTGGCGGCAA were annealed together and introduced into the Nhe I site thus constructing a vector (pCEP4/N -Strep) producing an N-terminal Strep II fusion protein.

Production of expression constructs for the expression of TGx.

Two episomal expression constructs were built allowing the production of both a C-terminal and an N-terminal Strep II fusion protein. For the construct with C-terminal Strep II tag, the TGx cDNA was amplified by polymerase chain reaction (PCR) using the 5'-prim r 5'-ATTGCGGCCGCCATG-GCCCAAGGGCTAGAAG, and the 3'-primer 5'-TAAGCGGCCGCTAATGCAAAGTCTACATAAAC. The 5'-primer introduced a new Not I restriction site and a Kozak's translation initiation sequ nce,

while the 3'-primer inserted a new Not I restriction site and removed the stop codon. The Not I restriction enzyme digested PCR product was purified and inserted at the same restriction sites of pCEP4 in order to obtain the final expression vector pCEP4/TGx/C-Strep. For the construct with N-terminal Strep II tag, the TGx cDNA was amplified by polymerase chain reaction (PCR) using the 5'-primer 5'-ATTGCTAGCCCAAGGGCTAGAAGTGG, and the 3'-primer 5'-TAAGCGGCCGCTTATA-ATGCAAAGTCTACATAAAC. The 5'-primer introduced a new Nhe I restriction site and removed the first methionine, the 3'-primer inserted a new Not I restriction site directly after the stop codon. After digesting with the restriction enzymes Nhe I and Not I the PCR product was purified and inserted at the same restriction sites of pCEP4 in order to obtain the final expression vector pCEP4/N-Strep/TGx.

Construction of expression constructs for the expression of TGk.

Two episomal expression constructs were built allowing the production of both a C-terminal and an N-terminal Strep II fusion protein. For the construct with C-terminal Strep II tag, the TGk proenzyme cDNA was amplified by polymerase chain reaction (PCR) using the 5'-primer 5'-ATTAAGCTTGCCGCCACCATGATGGATGGGCCACGTTCC, and the 3'-primer 5'-ATTGCGGCCGCGAGCTCCACCTCGAGATGCCATAGG. The 5'-primer introduced a new Hind III restriction site and a Kozak's translation initiation sequence, while the 3'-primer inserted a new Not I restriction site and removed the stop codon. The Hind III/Not I restriction enzyme digested PCR product was purified and inserted at the same restriction sites of pCEP-Pu/BM40SP/C-Strep, in order to obtain the final expression vector pCEP-PuTGk/C-Strep. For the construct with N-terminal Strep II tag, the TGk proenzyme cDNA was amplified by polymerase chain reaction (PCR) using the 5'-primer 5'-ATTGCTAGCAGATGGGCCACGTTCCGATG, and the 3'-primer 5'-ATTGGATCCTAAGCTCCACCTCGAGATGC. The 5'-primer introduced a new Nhe I restriction site and removed the first two methionines, while the 3'-primer inserted a new Not I restriction site directly after the stop codon. After digesting with the restriction enzymes Nhe I and Not I the PCR product was purified and inserted at the same restriction sites of pCEP4, in order to obtain the final expression vector pCEP4/N-Strep/TGk. The correct insertion and sequence of the full constructs was verified by cycle sequencing as described above.

Recombinant expression of the human TGe, TGx, and TGk.

Human embryonic kidney cells (293-EBNA, Invitrogen) were transfected and harvested in cell culture in Dulbecco's MEM NUT MIX F-12 (Life Technologies) medium containing 10% foetal bovine serum (Life Technologies), 1% L-glutamine (Life Technologies), 200 IU/ml penicillin (Life Technologies), and 200 µg/ml streptomycin (Life Technologies). Cells transfected with pCEP-Pu constructs were selected with 0.5 µg/ml puromycin (Sigma), those with pCEP4 with 333 µg/ml (335 U/ml) hygromycin B (Calbiochem). After removing the medium and washing with cold (4°C) 0.25 M sucrose, the cells were lysed mechanically in cold 0.25 M sucrose. The lysate was cleared of

particulate material by centrifugation at 27,200 g for 30 min at 4°C, followed by ultracentrifugation of the supernatant at 210,000 g for 60 min at 4°C. After filtering with cheesecloth and adding 1 mM PMSF (Fluka) as proteinase inhibitor, 12-72 ml of the supernatant was passed over a StrepTactin™ affinity column of 3 cm³ volume equilibrated with sterile and filtered 100 mM Tris/HCl, pH 7.5, containing 1 mM EDTA, at 4°C, and at a flow rate of 0.4 ml/cm²/min. After extensive washing with equilibration buffer containing 1 mM PMSF, at a flow rate of 0.9 ml/cm²/min, the protein was eluted with equilibration buffer containing 1 mM PMSF and 2.5 mM desthiobiotin (Sigma), at a flow rate of 0.4 ml/cm²/min. 2 ml fractions were collected. The purification was controlled by SDS-PAGE and immunoblotting with monoclonal antibodies against the Strep tag as described above. The protein concentration was estimated by SDS-PAGE, and measured using the bicinchoninic acid (BCA) protein assay reagent (Pierce) following the protocol provided by the supplier, with bovine serum albumin as a standard.

Transglutaminase activity assay

TGc and TGe activity was measured by incorporation of [1,4-³H]putrescine (Amersham) 30 min. at 37°C as described before (Aeschlimann D et al., J Biol Chem 1991;266:15308-17), with the only difference that the buffer contained 7.5 mM dithiothreitol in order to reduce any oxidised sulfhydryl groups important for catalytic activity. The TGe was activated by partial proteolytic digestion preincubating it 20 min at 37°C together with either 45.4 µg/ml (0.5 U/ml) proteinase K (Sigma), or 45.4 µg/ml (55.4 U/ml) trypsin 1:250 (Sigma), or 1.18 mg/ml (1 U/ml) dispase (Life Technologies).

Mass spectrometry

Mass spectrometry of TGc and TGe was performed by matrix-assisted laser desorption method using a Bruker Reflex III instrument equipped with a high mass detector for linear detection. Sinapinic acid was used as matrix, and external calibration was carried out using singly, doubly and triply charged molecular ions of protein A.

Sera and patients

In the first study, the patients had been examined at the Gastroenterological Departments of Internal Medicine or Paediatrics and the Department of Dermato-Venereology of the Semmelweis University. The CD diagnosis was confirmed by jejunal biopsy while DH was proven by skin biopsy. Serum samples were taken from 71 patients with GSE (33 with DH, and 38 with CD), 26 with non-CD gastrointestinal diseases (such as M. Crohn, food hypersensitivity, food intolerance, intestinal infection, reflux oesophagitis, non-CD diarrhoea, alimentary dystrophy), and 27 with other diagnoses like autoimmune diseases (systemic lupus erythematosus, diabetes mellitus type I), different skin disorders (pemphigus foliaceus, ichthyosis, urticaria), cholelithiasis, hepatosplenomegalia, nanosomia, and healthy controls. The mean age and sex of patients' groups are presented in Table 1.

TABLE 1
Age (at the time of blood sampling, in years) and sex of patients

	CD	DH	GI	Others	Altogether
Mean age	18.5	30.2	17.4	17.0	20.8
Min. age	3	6	1	12	1
Max. age	51	74	78	53	78
Male/female	20/18	13/20	14/12	14/13	61/63

5 To obtain data on the sensitivity of TGc ELISA, we included in the current study sera from 16 treated patients (patients on a gluten-free diet).

In the supplementary study on GSE associated autoimmune diseases the sera were obtained from the Gastroenterological Department of Internal Medicine or Paediatrics and the Department of Dermato-Venereology of the Semmelweis University, the Departments of the Internal Medicine I-IV of
10 the Medical Faculty of the University Cologne, and from the Laboratory for Autoimmune Diseases of the Wieslab Co. Sweden. Serum samples were taken from patients with the following autoimmune diagnoses (number of patients in parentheses): GSE (141; from which 73 DH including 18 on a gluten-free diet, and 68 CD including 27 on a gluten-free diet), Crohn disease (31), bullous pemphigoid (44), pemphigus vulgaris (57), colitis ulcerosa (21), Goodpasture syndrome (20),
15 Wegener granulomatosis (20), rheumatoid arthritis (44), SLE (25), progressive systemic sclerosis (7). Sera from patients with psoriatic arthritis (5) and hepatitis C (39) were also studied as these diseases may have an autoimmune component.

All serum samples were stored at -78 C until assayed. Among the control sera (48) some were from healthy individuals (21) and some from patients suffering from disease that does clearly not
20 have an autoimmune component (27). All serum samples were stored at -78 C until assayed.

EMA test.

Serum IgA antibodies were measured by an indirect immunofluorescence method (Collin P et al, Scand J Gastroenterol 1992;27:367-71). All serum samples were diluted 1:5 in phosphate-buffered
25 saline (PBS, pH: 7.4). 10 µm cryostat tissue slides of the aboral part of monkey (Cercopithecinae family) oesophagus were used as antigen. Bound IgA was detected by α-chain specific, fluorescein isothiocyanate-conjugated rabbit anti-human IgA antibodies (1:40 in PBS; Dako). All sera used in the first study were indisputably negative or positive for IgA EMA. In the supplementary study, no EMA test was performed.

ELISA.

The ELISA method was similar to the calcium-activated test described previously (Dieterich W et al, Gastroenterology 1998;115:1317-21; Sulkanen S et al., Gastroenterology 1998;115:1322-8). Briefly: 96 well microtiter plates (Nunc MaxiSorp) were coated by 1 µg guinea pig TGc (Sigma) or human TGe or TGc per well in 100 µl of 50 mM Tris/HCl, pH 7.5, containing 5 mM CaCl₂ per well at 4°C overnight (at least 9 hours). No blocking was used. After each step the wells were washed by 50 mM Tris/HCl containing 10 mM EDTA and 0.1% Tween 20 (TET). Sera were diluted to various concentrations with TET, or preincubated in 50 mM Tris/HCl containing 0.1% Tween 20 and various concentrations of TGc or TGe or guinea pig TG, and incubated on the plates for 1.5 hours at room temperature. Bound IgA was detected by peroxidase-conjugated anti-human IgA antibody (Dako), diluted 1:4000 in TET and incubated for 1 hour at room temperature. The colour was developed by 100 µl of 60 µg/ml 3,3',5,5'-tetramethylbenzidine substrate in 100 mM sodium acetate, pH 6.0, containing 0.015% H₂O₂ for 5 minutes at room temperature. The reaction was stopped by adding 100 µl of 20% H₂SO₄. The absorbance was read in an ELISA reader at 450 nm.

The amount of protein and the serum concentrations used in the test were optimised. All serum samples were examined in triplicates, and triplicates of a negative and a positive reference serum, as well as a buffer blank were included in each plate. The antibody concentrations were expressed in arbitrary units (AU), i.e. as percentages of the positive reference serum.

In order to get data about the effects of calcium-activation, an experiment coating with human TGc without CaCl₂ in the coating buffer was also performed.

Inhibition ELISA.

Serum IgA levels against TGe were compared with those against TGc. Four sera from patients with CD or DH were studied which had previously shown to contain slightly (1 serum), moderately (1 serum) or highly (2 sera) elevated serum IgA antibody titers against TGc. Blocking experiments were carried out by preincubating the sera at a serial dilution from 1:250 to 1:32000 in 50 mM Tris/HCl containing 0.1% Tween 20 and 1 µg of either TGe or TGc 60 min at room temperature. A serum dilution was chosen at which the greatest difference was detected between the antibody titers of the sera with and without preincubation. In the following inhibition experiment the four sera were preincubated at this dilution in 50 mM Tris/HCl containing 0.1% Tween 20 with a serial dilution ranging from 0-4 µg of either TGc or TGe 60 min at room temperature.

Statistics

Optical densities (and thus titres given in AU values) did not show Gaussian distribution, thus for statistical description of titers from the different patient groups, we present medians with their 95% confidence intervals (95% CI) (Gardner MJ & Altman DG, eds. Statistics with confidence-confidence intervals and statistical guidelines. London: British Medical Journal, 1989:28 pp.) and for comparison

Mann-Whitney's non-parametric, unpaired, two-tailed test was used (Werner J. Biomathematik und Medizinische Statistik, 2nd ed. München-Wien-Baltimore: Urban & Schwarzenberg, 1992:53 pp.). For describing correlation of titers, Spearman's correlation coefficient with its 95% CI and correlation analysis for unpaired data of non-normal distribution was used (Gardner and Altman, 1989; Werner, 1992).

For comparison of titres in the calcium-activated and inactivated human TGc ELISA, Wilcoxon's two-tailed signed rank test was performed (Werner, 1992). For describing and comparing of the two ELISA systems, the receiver operating characteristic (ROC) curves and the areas under the ROC curves (AUC) with their 95% CI are presented. For calculating confidence intervals of AUC, beside the most often used method also a bootstrap technique, the bias-corrected and accelerated (BC_a) confidence interval method was applied, as it is more appropriate in describing confidence intervals of AUC which are very close to the maximum (1.0) (DeLong ER et al. Biometrics 1988;44:837-45; Mossman D., Med Decis Making 1995;15:358-66; Hellmich M. Receiver operating characteristic (ROC) Kurven und Flächen darunter. [PhD thesis] 1996. <http://www.medizin.uni-koeln.de/kai/imsie/homepages/Martin.Hellmich/dr.html> (accessed June 1999).

EXAMPLE

Recombinant human TGc.

The human TGc was expressed in the 293-EBNA human embryonic kidney cell line as a fusion protein with the Strep II tag. The protein could be purified in a single step by affinity binding to a StrepTactin column, on washing with desthiobiotin the protein eluted as a single band with an estimated molecular weight of 89 kDa (Fig. 1A) when visualised by Coomassie-stained SDS-PAGE. Western blot analysis showed that the band reacted with monoclonal anti-TGc antibodies (Fig. 1B). The column bound almost all the tagged protein with no immunoreactivity appearing in the flow through (Fig. 1B). The yield from the lysate of a confluent cell monolayer in a cell culture dish of 13 cm diameter was approximately 200 µg. The molecular mass calculated from the sequence of the human TGc is 77.3 kDa, and the calculated molecular mass of the fusion protein (TGc having a carboxyterminal tag of 10 amino acids) is 78.4 kDa. Mass spectrometry of the fusion protein gave a molecular mass of 78.3 kDa. In cell lysates, the activity of the expressed human TGc was 4.7 times higher than the background activity of transglutaminases present in untransfected 293-EBNA cells. The freshly purified human TGc showed similar or higher activity than the guinea pig TGc from Sigma.

Performance of the human TGc ELISA.

The optimal coating concentration of human TGc was 1 µg per well. Using highly positive sera from four patients for calibration, a log-linear curve was seen between dilutions of 1:250 and

1:32000. Four negative sera showed some signal at lower dilutions ($>1:500$). Some positive sera showed a signal plateau at dilutions of 1:250 or less. The ratio between the mean OD values of positive and negative results at the dilution of 1:125 was 1:6, while at greater dilutions more than 1:10. Hence in the assay a serum dilution of 1:250 was used. One positive and one negative reference serum sample was included in each assay to control the test performance. The positive serum was used as "standard", and the optical density results were given as arbitrary units (AU) calculated as a percentage of the standard serum. The mean intra- and interassay coefficients of variation for the standard serum were 1.3% and 13.7%, respectively. The mean intra- and interassay coefficients of variation for human TGc ELISA were 3.2% ($n=124$) and 9.2% ($n=15$), respectively.

The median antibody concentrations for the patients with untreated GSE (CD or DH) was 61.4 AU ($n=55$, 95% CI: 45.1-78.5), for controls 12 AU ($n=53$, 95% CI: 10.8-13), the difference was significant ($p<0.0001$). For treated patients, the median of antibody concentrations was 48.1 AU ($n=16$, 95% CI: 20.8-85.6), for controls with gastrointestinal diseases 12.1 AU ($n=26$, 95% CI: 9.8-14.7), for healthy individuals and controls with other diagnoses 12 AU ($n=27$, 95% CI: 10.7-13.0), respectively. The area under the ROC curve was 0.999 (95% CI: 0.996-1.001; 95% CI with BC_a method: 0.990-1.0) (Fig. 2).

A cut-off value of 18 AU was chosen, and sera with antibody concentrations equal or higher than 18 AU were labelled as human TGc ELISA positive. This cut-off value gave a specificity and a sensitivity of 98.1% (95% CI: 95.7-100%) and 98.2% (95% CI: 95.9-100%), respectively (treated patients are excluded). The coincidence of the human TGc assay with the clinical diagnosis (excluding treated patients) was 106/108 (98.1%), giving one false positive and one false negative result (Fig. 3).

Recombinant human TGe

The human TGe was expressed in the 293-EBNA human embryonic kidney cell line as a fusion proenzyme with the Strep II tag. The protein could be purified in a single step by affinity binding to a StrepTactinTM column. After elution with desthiobiotin the protein gave a single 80 kDa band in SDS-PAGE (Fig. 9), which reacted with monoclonal antibodies against the Strep II tag (Fig. 10, lane II). The column bound almost all the tagged TGe with no immunoreactivity appearing in the flow through. The yield from the lysate of a confluent cell monolayer in a cell culture dish of 13 cm diameter was approximately 200 μ g. The molecular mass calculated from the sequence of the human TGe proenzyme is 76826 Da, and the calculated molecular mass of the fusion protein (TGe proenzyme having a carboxyterminal tag of 10 amino acids) is 78011 Da. Mass spectrometry of the fusion protein gave a molecular mass of 77765 Da. In cell lysates, the activity of the expressed human TGe was 2.5 times higher than the background activity of transglutaminases present in untransfected 293-EBNA cells. The freshly purified human TGe proenzyme showed 1;40-1;80 of the

activity of TGc and when activated with different proteases (proteinase K, trypsin or dispase) similar or higher activity than the TGc.

Performance of human TGe ELISA.

5 GSE sera showing elevated IgA antibody levels against the human TGc also showed elevated IgA titres against the human TGe, whereas sera not reacting with the human TGc did not react with the human TGe (Fig. 11). The antibody titres for TGe were in most cases lower than for TGc (Fig. 11).

10 Inhibition ELISA.

Preincubation of GSE sera with human TGc or TGe inhibited the IgA reactivity in an ELISA plate coated with human TGc (Fig. 12). The results shown support that serum IgA antibodies from patients with CD and DH react with both the human TGc and the TGe, although the titres to TGe is lower. Both TGc and TGe can inhibit the reaction of serum IgA antibodies with TGc. At least a part of the serum antibodies from patients with CD and DH is directed against epitopes which are shared by the two transglutaminases.

Performance of the guinea pig TGc ELISA.

20 The optimal coating concentration of guinea pig TGc was 1 µg per well and the optimal serum dilution 1:250, as with the human TGc ELISA. Each assay was performed parallel to the human TGc assay at the same time, and the same serum samples and serum dilutions were used. The mean intra- and interassay coefficients of variation of the standard serum were 2.2% and 9.0%, respectively. The intra- and interassay coefficients of variation for guinea pig ELISA were 2.8% (n=124) and 12.8% (n=15), respectively.

25 The median of antibody concentrations for the patients with untreated GSE (CD or DH) was 51.8 AU (n=55, 95% CI: 34.2-63), for controls 8 AU (n=53, 95% CI: 7.3-8.9), the difference was significant (P<0.0001). For treated patients, the median of antibody concentrations was 18 AU (n=16, 95% CI: 9.2-69.9), for controls with gastrointestinal diseases 7.5 AU (n=26, 95% CI: 6.6-9), for healthy individuals and controls with other diagnoses 8.5 AU (n=27, 95% CI: 7.2-10.3), respectively. The area under the ROC curve was 0.980 (95% CI: 0.958-1.002; 95% CI with BC_a method: 0.943-0.993) (Fig. 4).

30 A cut-off value of 14 AU was chosen, and sera with antibody concentrations equal or higher than 14 AU were labelled as guinea pig TGc ELISA positive. This cut-off value gave (excluding treated patients) a specificity and a sensitivity of 96.2% (95% CI: 92.8-99.6%) and 92.7% (95% CI: 88.1-97.3%), respectively. The coincidence of the guinea pig TGc assay with the clinical diagnosis (excluding treated patients) was 102/108 (94.4%), giving 2 false positive and 4 false negative results (Fig. 5).

Effects of calcium-activation.

32 serum samples were tested in ELISA for IgA anti-human TGc antibodies with and without calcium-activation. The overall antibody titres did not show significant difference ($p=0.27$). However, sera with anti-TGc titres lower than 30 AU in the calcium-activated assay were significantly lower in the assay without calcium-activation ($n=18$, $p=0.009$), whereas higher titres were not significantly different ($n=14$, $p=0.35$).

Comparison of EMA with TGc ELISA.

Excluding treated patients, with the exception of one false positive result, all patients with EMA positive sera had GSE (55/56, 98.2%). 12 of 16 (75%) treated patients with GSE were positive for EMA. Comparing only the untreated EMA positive cases, the results of human and guinea pig TGc ELISA coincided with the EMA test in 54/56 (96.4%) and 51/56 (91.1%) cases, respectively. The false positive serum by EMA was negative by both human and guinea pig TGc ELISAs. The one false negative serum with human TGc ELISA was also negative with guinea pig TGc ELISA. The 12 patients on incomplete gluten-free diet with EMA positivity had also positive anti-TGc IgA titres with both ELISA systems.

All patients negative for EMA were either treated patients having GSE or patients not having GSE. Comparing only the untreated EMA negative cases, the results of human and guinea pig TGc ELISA coincided with the EMA test in 52/53 (98.1%) and 51/53 (96.2%) cases, respectively. The one false positive serum with human TGc ELISA was also positive with guinea pig TGc ELISA; in addition, another false positive serum was detected by the guinea pig assay. Both false positive sera were from patients with M. Crohn. The four EMA negative patients with treated GSE were also negative by guinea pig TGc ELISA, but one of them was positive by human TGc ELISA.

The overall coincidence of EMA test with human and guinea pig ELISA was 120/124 (96.8%) and 117/124 (94.4%), respectively.

Comparison of human TGc ELISA with guinea pig TGc ELISA.

The results of the two ELISAs coincided in 119/124 (96%) of all tested sera. In four discordant cases, the human assay was more sensitive than the guinea pig assay, giving positivity in the human assay. One of them was an EMA negative CD patient on gluten-free diet. In the fifth discordant case, the guinea pig ELISA gave false positive result for a patient with M. Crohn. The antibody titre was, however, also high (17.5 AU) in the human TGc ELISA, almost reaching the cut-off level (18 AU).

The false results of the human TGc ELISA coincided with those of the guinea pig TGc ELISA. Both assays failed to recognise the serum of one EMA positive CD patient, and both detected

a patient having Crohn's disease as positive. Both tests gave correct, negative result in the case of a false EMA positive patient.

The titres with the two assay correlated well ($r_s=0.9377$, 95% CI: 0.9121-0.9559, $p<0.0001$), the correlation was theoretically exponential, but in the practice linear, with an exponent of 1.05 (Fig. 8). The difference between the areas under the ROC curves was 0.019 (95% CI: -0.002-0.040; 95% CI with BC_a method: 0.005-0.056).

Recombinant human TGe

The human TGe was expressed in the 293-EBNA human embryonic kidney cell line as a fusion proenzyme with the Strep II tag. The protein could be purified in a single step by affinity binding to a StrepTactin™ column. After elution with desthiobiotin the protein gave a single 80 kDa band in SDS-PAGE (Fig. 9), which reacted with monoclonal antibodies against the Strep II tag (Fig. 10). The column bound almost all the tagged TGe with no immunoreactivity appearing in the flow through. The yield from the lysate of a confluent cell monolayer in a cell culture dish of 13 cm diameter was approximately 200 µg. The molecular mass calculated from the sequence of the human TGe proenzyme is 76826 Da, and the calculated molecular mass of the fusion protein (TGe proenzyme having a carboxyterminal tag of 10 amino acids) is 78011 Da. Mass spectrometry of the fusion protein gave a molecular mass of 77765 Da. In cell lysates, the activity of the expressed human TGe was 2.5 times higher than the background activity of transglutaminases present in untransfected 293-EBNA cells. The freshly purified human TGe proenzyme showed 1;40-1;80 of the activity of TGc and when activated with different proteases (proteinase K, trypsin or dispase) similar or higher activity than the TGc.

Discussion of Example and Results

To our knowledge, human TGc had not been expressed before in mammalian cells via DANN recombination methods. We preferred using human cells instead of bacteria for two reasons. First, although there is no evidence for posttranslational modifications of TGc, we can not completely exclude this possibility. Such modifications would in all probability not occur in bacteria. Second, if chaperons are needed to obtain a correct folding, these are more likely to be present in human cells.

The molecular mass of the guinea pig TGc differs only slightly (0.1 kDa) from that of the human TGc when measured by mass spectrometry, but the guinea pig TGc runs appreciably faster on SDS-PAGE than the human TGc. The TGc from human fibroblasts runs with the same speed as our fusion protein, although the difference between them is 1.2 kDa. These observations imply that the difference between the structures of the human and the guinea pig TGc is more profound than suggested by their high amino acid identity.

The human TGc as a fusion protein with the Strep II tag could be purified very effectively in one step giving one single band on a Coomassie-stained gel. The protein band could be clearly identified by immunoblot as tissue transglutaminase. Mass spectrometry gave the expected molecular size. The purified recombinant human protein had transglutaminase activity. It is not known whether the human TGc has the same catalytic activity as the guinea pig enzyme. Our purified protein had similar or higher activity as the same amount of guinea pig liver TGc. Thus we conclude that the protein we used in the assays is the purified, active, human TGc with a C-terminal Strep II tag.

The guinea pig liver TGc preparation used for testing contains other protein contaminants which are not immunoreactive with monoclonal anti-TGc antibodies. However, as it had been used successfully by other authors in its original form, we did not purify it further. It can not be excluded that immunopositivity seen in microtiter wells might in some cases be due to reactivity against contaminants.

Sulkanen and co-workers (Sulkanen S et al. Gastroenterology 1998;115:1322-8.) demonstrate that calcium given to the coating buffer, in a concentration suitable for activation of guinea pig TGc, causes improved test performance by more specific detection of TGc by antibodies. The authors explain their observation as a conformational change occurring through calcium-activation, but after coating they remove calcium by washing and as they then use a high concentration of EDTA, so the antibody-antigen reaction occurs in the absence of calcium. Thus a conformational change is assumed to be retained by the immobilised TGc molecules after removal of calcium. Besides conformational changes, which are unlikely to become irreversible by coating, there is another possible explanation for this effect. TGc can act as its own substrate, and is able to cross-link itself to form protein complexes of high molecular weight (Birckbichler PJ et al, Biochem Biophys Res Commun 1977;78:1-7). This is likely to occur in the wells where the concentration of TGc is approximately the same as used in the activity assay. Thus not only TGc monomers, but also TGc complexes may be immobilised in the ELISA wells and this may favour the binding of antibodies giving a higher signal. In addition, cross-linked TGc complexes may reveal new antigenic epitopes, different from those of the TGc monomer. This hypothesis is supported by the observation that preincubation of CD sera with calcium-activated guinea pig TGc blocks effectively the reactivity in ELISA, but to obtain the same blocking effect with inactivated guinea pig TGc, a preincubation with a tenfold amount is needed.

Sulkanen and co-workers (Sulkanen S et al., Gastroenterology 1998;115:1322-8) changed the original method (Dieterich W et al, Nature Medicine 1997;3(7):797-801) not only by using calcium activation, but also changed the buffer and omitted blocking. They do not present data showing which component is responsible for the improved assay performance. We tested some positive and

negative sera with and without calcium-activation of the human TGc, and could not find significant changes in the overall performance of the test. This suggests that other factors, in addition to calcium, may be important in the optimisation of the guinea pig TGc assay. However, as the optical densities of EMA positive sera with anti-TGc titres lower than 30 AU in the calcium-activated assay were significantly lower, or even negative, in the assay without calcium-activation, we use calcium also in the assay with human TGc.

The setting of the cut-off values for the ELISAs was based on the ROC-analysis of the tests. A cut-off value providing perfect separation of individuals having or not having GSE could not be found, though the coincidence with the diagnosis by biopsy is in both tests very high. The guinea pig TGc ELISA was not able to detect two untreated patients with DH and one untreated patient with CD which could be detected by the human TGc ELISA. The 95% confidence intervals of sensitivities of the two ELISAs overlap, and therefore the sensitivity difference has to be confirmed by further studies, but the results affirm the assumption that in a few cases autoantibodies are directed against epitopes of human TGc not conserved in guinea pig TGc. One serum from a CD patient with clear EMA positivity was not immunoreactive in either ELISA, and the titre values were so far below the cut-off that the result is probably not due to chance. It is conceivable that TGc is not the only autoantigen in GSE, and that in some rare cases, even though EMA positivity occurs, no antibodies against TGc are present. This is supported also by the observation that the immunoabsorption of IgA-class autoantibodies against TGc by guinea pig TGc cannot completely abrogate anti-endomysium activity (Lock RJ et al., Clin Exp Immunol 1999;116:258-62), although these experiments should be repeated using the calcium-activated human TGc, as not all patients' antibodies may cross-react with the guinea pig antigen.

The EMA test gave a false positive result in an 8-year-old girl who had a transient diarrhoea in February 1998. Repeated EMA tests showed IgA binding in the intercellular spaces of smooth muscle cells. Jejunal histology was negative for GSE, and the diarrhoea has not recurred. The fact that a serum was false positive by EMA test, but correctly diagnosed by both ELISAs, also underlines the possibility of EMA positivity due to antigens other than TGc.

It is interesting that two EMA-negative patients with Crohn's disease had TGc antibody titres above the cut-off level in the guinea pig TGc ELISA, one of them also in the human TGc ELISA. Coeliac disease and Crohn's disease have been described to occur in the same patient (Gillberg R et al, Scand J Gastroenterol 1982;17:491-6), but this association is very rare. In our two cases associated coeliac disease can not be ruled out, but as both titres are near to the borderline (21.6 and 17.5 AU in the human TGc ELISA, 15.9 and 17.8 AU in the guinea pig ELISA), the elevation of titres might result from a low level IgA autoantibody production against TGc in Crohn's disease rather than

from that in active coeliac disease. This speculation is supported by the fact that in both ELISA systems the median of titres of patients with Crohn's disease is greater than that of healthy individuals and patients with other gastrointestinal or non-gastrointestinal diseases. However, the differences and the number of patients' sera tested in the present study are too small to allow us to judge the significance of this finding.

Patients on complete or incomplete gluten-free diet had a wide spectrum of antibody titres, and the results of the ELISAs were in good agreement with those of EMA tests. The ELISA with the human antigen turned out to be slightly more sensitive in this regard than the EMA test or the guinea pig ELISA, recognising one EMA- and guinea pig ELISA negative CD patient as positive.

Compared to the other established systems, the human TGc ELISA was found to be as specific and sensitive as the EMA test, and somewhat superior to the guinea pig TGc ELISA. The results show the high diagnostic value of all tested systems in this study, but in particular that of the human TGc ELISA, which has almost perfect sensitivity and specificity, and does not have the disadvantages of EMA test. Thus we conclude that the human TGc-based ELISA should be the method of choice for easy and non-invasive screening and diagnosis of GSE.

The results summarised in Figures 6 and 7 show that IgA against human tissue transglutaminase is also detected in autoimmune disorders other than gluten sensitive enteropathy. The positivity in the human TGc ELISA among patients with autoimmune disease other than CD and DH is most probably not due to these patients having an active GSE. It is indeed very unlikely that more than the quarter of all autoimmune patients would have associated GSE, even though it is conceivable that in some cases where the titres are very high the test shows a real association. Celiac disease has been described to associate with various autoimmune disorders as shown in Table 2 below:

TABLE 2

Forms of autoimmune (AI) diseases that have been reported to associate with GSE:

Addison's disease	Recurrent pericarditis
AI haemolytic anaemia,	Relapsing polychondritis
AI thrombocytopenic purpura	Rheumatoid arthritis
AI thyroid diseases	Sarcoidosis
Atrophic gastritis - pernicious anaemia	Sjögren's syndrome
IgA nephropathy or IgA glomerulonephritis	SLE, splenic atrophy
Myasthenia gravis	Type I (insulin-dependent) diabetes mellitus
Partial lipodystrophy	Ulcerative colitis
Polymyositis	Vasculitis (both systemic and cutaneous)
Primary biliary cirrhosis	Vitiligo
Primary sclerosing cholangitis	

A part of these associations are proven, the other part have been anecdotal.

5

It has been known that GSE can provoke a T-cell mediated inflammation of the duodenojejunal bowel region causing various resorption disturbances (for review see Trejdosiewicz LK et al., Clin Gastr 1995;9:251-72). The latter may result in painful diarrhea, sideropenic anemia, hypoproteinemia, osteoporosis, amenorrhea, hypovitaminoses, and in children in retardation of growth and development (for review see Corazza GR, Gasbarrini G. Coeliac disease in adults. Bailliere Clin Gastr 1995;9:329-50; Littlewood JM. Coeliac disease in childhood. Bailliere Clin Gastr 1995;9:295-328.). Beside these direct consequences the persistence of the disease predisposes of various autoimmune disorders (e.g. diabetes type 1), and malignancies (e.g. duodenojejunal lymphomas). The clinical signs and symptoms of DH are mainly those of skin affection (polymorphic, itching blisters with underlying erythema typically located over the extensor surfaces of the big joints), the gastroenterological symptoms are often mild or clinically completely absent. However, the inflammatory small bowel changes can often be found by histological examination even if there are no clinical signs or symptoms suggesting jejunal pathology. The enteropathy in DH is morphologically, clinically and functionally identical with that in CD suggesting identical or very similar aetiology and pathomechanism of both of these two forms of GSE.

20

Our results however show a general role of TGc and other transglutaminases in autoimmune processes. As a consequence positive results for TGc IgA in patients suffering from other autoimmune disease should not alone be taken as the basis for the diagnosis of GSE. On the other hand, antibodies against TGc and other transglutaminases such as TGe can be used as a marker for

25

patients suffering from other autoimmune disease of the GSE-type and may serve to define subgroups of patients within such disease groups.


Consequently, we provided a new method for diagnosis of autoimmune diseases of the GSE-type or associated with gluten sensitive enteropathy, essentially comprising taking a specimen and testing the specimen for antibodies against human tissue transglutaminase, or other transglutaminases. In this way autoimmune diseases other than coeliac disease can be diagnosed and distinguished, notably, dermatitis herpetiformis Dühring, Addison's disease, AI haemolytic anaemia, AI thrombocytopenic purpura, AI thyroid diseases, atrophic gastritis - pernicious anaemia, IgA nephropathy or IgA glomerulonephritis, myasthenia gravis, partial lipodystrophy, polymyositis, primary biliary cirrhosis, primary sclerosing cholangitis, recurrent pericarditis, relapsing polychondritis, rheumatoid arthritis, rheumatism, sarcoidosis, Sjögren's syndrome, SLE, splenic atrophy, type I (insulin-dependent) diabetes mellitus, diabetes mellitus of other types, ulcerative colitis, vasculitis (both systemic and cutaneous), vitiligo as well as autoimmune diseases associated with female infertility (Collin et al, Gut, 1996: 39, 382-384), increased risk of abortion (Smecul et al., Eur. J. Gastr. & Hep. 1996; 8(1), 63-67), or reduced foetal growth due to the presence of an autoimmune disease of the GSE-type or latent, non-active GSE.

CLAIMS:

1. Method for diagnosis of autoimmune diseases of the GSE-type or associated with gluten sensitive enteropathy comprising taking a sample and testing the sample for antibodies against human tissue transglutaminase and at least one other transglutaminase molecule.
2. Method as claimed in claim 1, wherein the autoimmune disease is dermatitis herpetiformis, morbus Duhring.
3. Method as claimed in claim 1, wherein the autoimmune disease is selected from Addison's disease, AI haemolytic anaemia, AI thrombocytopenic purpura, AI thyroid diseases, atrophic gastritis - pernicious anaemia, Crohn's disease, colitis ulcerosa, Goodpasture syndrome, IgA nephropathy or IgA glomerulonephritis, myasthenia gravis, partial lipodystrophy, polymyositis, primary biliary cirrhosis, primary sclerosing cholangitis, progressive systemic sclerosis, recurrent pericarditis, relapsing polychondritis, rheumatoid arthritis, rheumatism, sarcoidosis, Sjögren's syndrome, SLE, splenic atrophy, type I (insulin-dependent) diabetes mellitus, diabetes mellitus of other types, Wegener granulomatosis, ulcerative colitis, vasculitis (both systemic and cutaneous), vitiligo.
4. Method as claimed in claim 1, wherein the autoimmune disease is associated with infertility, increased risk of abortion, reduced foetal growth.
5. Comparative protein binding assay for a differential diagnosis of autoimmune diseases comprising the detection of antibodies against transglutaminase, characterised in that the protein binding assay comprises recombinant human tissue transglutaminase as antigen.
6. Comparative protein binding assay as claimed in claim 5, further comprising tissue-specific transglutaminase as antigen.
7. Comparative protein binding assay as claimed in claim 6, wherein the transglutaminase is selected from FXIIIA, TGk, TGe, TGx, and band 4.2.
8. Comparative protein binding assay as claimed in any preceding claim 5 or 6, further comprising any other transglutaminase as antigen.

9. Comparative protein binding assay as claimed in claim 8, comprising transglutaminase from different species.
10. Comparative protein binding assay as claimed in claim 8, wherein the transglutaminase is a recombinant fusion protein or fragment thereof.
11. Comparative protein binding assay as claimed in any preceding claim 5 to 10, wherein the binding assay is an immunoassay selected from RIA, EIA/ELISA, LiA and FiA.
- 10 12. Comparative protein binding assay as claimed in any preceding claim 5 to 11, wherein the binding assay is a sandwich-immunoassay selected from IRMA, IEMA/EUA, ILMA (immunoluminescence assay) and IFMA (immunofluorescence assay).

1 / 12

**Fig. 1**

2 / 12






Fig. 2

3 / 12

**Fig. 3**

4 / 12

**Fig. 4**

5 / 12




Fig. 5

6 / 12





Fig. 6

7 / 12

Fig. 7



8 / 12





Fig. 8

94—
67—
43—
30—


Fig. 9

10 / 12

**Fig. 10**

11 / 12

Fig. 11



12 / 12




Fig. 12